Seat No. : \_\_\_\_\_

# **JA-105**

### January-2021

## B.Sc., Sem.-III

## 201 : Mathematics

## (Advanced Calculus – I)

### Time : 2 Hours]

#### [Max. Marks : 50

**Instructions :** (1) Attempt any **three** questions from **Q-1** to **Q-8**.

- (2) **Q-9** is compulsory.
- (3) Notations are usual, everywhere
- (4) Figures to the right indicate marks of the question/sub-question.

1. (a) Define limit of function of two variables. Use the definition to find

$$\lim_{(x,y)\to(1,1)} \frac{x^2 + y^2}{xy}.$$

(1) 
$$f(x, y) = \begin{cases} \frac{x^2 - y^2}{x + y} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$
 at point (0, 0)

(2) 
$$f(x, y) = \begin{cases} \tan^{-1} \begin{pmatrix} y \\ x \end{pmatrix} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \text{ at point } (0, 1)$$

2. (a) Define iterated limits. Find the iterated limit for

$$f(x, y) = \begin{cases} \frac{x^2 - y^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 2 & \text{if } (x, y) = (0, 0) \end{cases} \text{ at point } (0, 0)$$

**JA-105** 

7

(b) Evaluate the following limit if exists

(1) 
$$\lim_{(x,y)\to(0,0)} f(x, y) \text{ where}$$
$$f(x, y) = \begin{cases} \frac{x^3 + y^3}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 2 & \text{if } (x, y) = (0, 0) \end{cases}$$

(2) 
$$\lim_{(x,y) \to (0,0)} f(x, y)$$
 where

$$f(x, y) = \begin{cases} \frac{\sin(x+y)}{x+y} & \text{if } (x, y) \neq (0, 0) \\ 1 & \text{if } (x, y) = (0, 0) \end{cases}$$

3. (a) State and prove Young's theorem.

(b) Find 
$$f_{xx}(0, 0)$$
,  $f_{yy}(0, 0)$ ,  $f_{yx}(0, 0)$  and  $f_{xy}(0, 0)$  for the function  

$$f(x, y) = \begin{cases} \frac{xy(x^2 - y^2)}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

### 4. (a) State and prove Schartz's theorem.

(b) Discuss the differentiability of the following functions :

(1) 
$$f(x, y) = \begin{cases} \frac{x^3 y^3}{(x^2 + y^2)^3} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$
 at point (0, 0)

(2) 
$$f(x, y) = x^2 + y^2$$
 at point (0, 0)

5. (a) If  $u = \phi(H)$  is function of a homogenous function H = f(x, y) of degree m whose partial derivatives of second order exists, then 7

(1) 
$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = m \frac{F(u)}{F'(u)} F'(u) (\neq 0) = G(u) \text{ (say)}$$

(2) 
$$x^2 \frac{\partial^2 u}{\partial x^2} + 2 xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = G(u) (G'(u) - 1)$$

Where H = f(x, y) = F(u).

**JA-105** 

7

7

7

7

2

	(b)	(1) If $f(x, y) = \sqrt{x^2 - xy}$ , then prove that	7
		$x^{2} \frac{\partial^{2} \mathbf{f}}{\partial x^{2}} + 2xy \frac{\partial^{2} \mathbf{f}}{\partial x \partial y} + y^{2} \frac{\partial^{2} \mathbf{f}}{\partial y^{2}} = 0$	
		(2) Find the extreme values of $f(x, y) = x^3 + y^3 - 3axy$ .	
6.	(a)	State and prove Euler's theorem for homogenous function.	7
	(b)	If $\mathbf{u} = f\left(\frac{\mathbf{y} - \mathbf{x}}{\mathbf{xy}}, \frac{\mathbf{z} - \mathbf{x}}{\mathbf{xz}}\right)$ , then prove that $x^2 \frac{\partial \mathbf{u}}{\partial x} + y^2 \frac{\partial \mathbf{u}}{\partial y} + z^2 \frac{\partial \mathbf{u}}{\partial z} = 0$ .	7
7.	(a)	Find the radius of curvature of a curve $r = f(\theta)$ i.e. in polar equations.	7
	(b)	State and prove Taylor's theorem for the function of two variables.	7
8.	(a)	(1) Expand $f(x, y) = e^{ax} \sin by$ in the power of x and y.	7
		(2) Find the radius of curvature of a curve $x^2 + y^2 = a^2$ .	
	(b)	Find the radius of curvature of parabola $r = a(1 - \cos \theta)$ .	7
9.	Attempt any <b>four</b> in short :		8
	(a)	Define multiple point and double point.	
	(b)	Define conjugate point.	
	(c)	Define harmonic function.	
	(d)	If $u = e^{xy}$ , then find $\frac{\partial^2 u}{\partial x \partial y}$ .	

(e) Find the degree of the homogenous function  $z = \frac{x^{21/3} + x^{7/2} y^{7/2}}{x^5 + y^5}$ .

(f) Give one example of function of two variables which is discontinuous at point (1, 1).

JA-105