Seat No.:	

## **MF-101**

## **March-2019**

## B.B.A., Sem.-III

**CC-206**: Elementary Statistics

7
61 11
C 1 11
on of ball and 2% is this and it roduced in
at random. l. Find her
1 a special tes, 39 are les person
4
bability of
·
11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

- 2. (A) (1) Obtain the probability mass function of Binomial distribution and state its properties.
  - (2) The number of accidents in a year attributed to taxi drivers in a city follows Poisson distribution with mean 3. Out of 1,000 taxi drivers, find approximately the number of drivers with (a) no accident in a year (b) more than 3 accidents in a year.  $(e^{-3} = 0.0498)$

7

7

7

7

4

7

7

OR

- (1) Assuming the probability of a female birth as  $\frac{1}{2}$ , find the probability that a family of 3 children will have (a) At least one boy (b) At most two boys.
- (2) There are 12 Indica and 8 Zen cars with a company. From them 5 cars are in repair in a workshop. Find the probabilities that of these cars: (a) there are 3 Indica and 2 Zen, (b) at least 3 Zen cars, (c) all the cars are of same type.
- (B) Answer the following: (Any Four)
  - (1) In distribution, Mean = Variance.
  - (2) The number of defects in a radio set is an example of Binomial distribution.

    (True/False)
  - (3) Write the probability mass function of Hyper Geometric distribution.
  - (4) Write the formula for finding mean of Hyper Geometric Distribution.
  - (5) The mean of Poisson distribution is 1.44, its S.D. = \_\_\_\_\_.
  - (6) When  $n \to \infty$  and  $P \to 0$  binomial distribution tends to Poisson distribution.
- 3. (A) (1) Define regression co-efficient. Give the properties of regression co-efficient.
  - (2) Calculate the co-efficient of correlation by Karl Pearson's method from the following data:

x	100	200	300	400	500	600	700
y	30	50	60	80	100	110	130

OR

MF-101 2

7

The following values have been obtained from the measurement of three (2) variables  $x_1$ ,  $x_2$  and  $x_3$ .

$$\bar{x}_1 = 6.8$$
  $\bar{x}_2 = 7.0$   $\bar{x}_3 = 7.4$ 

$$\bar{x}_2 = 7.0$$

$$\bar{x}_3 = 7.4$$

$$S_1 = 1.0$$

$$S_2 = 0.80$$

$$S_1 = 1.0$$
  $S_2 = 0.80$   $S_3 = 0.90$ 

$$r_{12} = 0.60$$

$$r_{12} = 0.60$$
  $r_{13} = 0.70$   $r_{23} = 0.65$ 

$$r_{23} = 0.63$$

Obtain the regression equation of  $x_1$  on  $x_2$  and  $x_3$ .

(B) Answer the following: (Any **Three**)

3

- (1) Define partial correlation.
- (2) The value of correlation co-efficient is between \_\_\_\_\_ and \_\_\_\_.
- The two regression co-efficients are 0.6 and 0.2, hence correlation co-(3) efficient is \_\_\_\_\_.
- Write a formula for  $r_{13.2}$ . (4)
- For a bivariate sample n = 16, r = 0.8, find probable error of r. (5)
- 4. Write a short note on Theory of Runs: (A) (1)

7

7

12 samples each of size 6 are drawn from a production process. Draw  $\overline{X}$ (2) and R chart. Also give your interpretations.

Sample No.	1	2	3	4	5	6	7	8	9	10	11	12
$\bar{\mathbf{x}}$	15	17	15	18	17	16	15	14	13	12	15	13
R	7	3	7	3	2	1	5	7	8	10	2	7

$$(n = 6, A_2 = 0.483, D_3 = 0, D_4 = 2.004)$$

OR

	(1)	Bajaj electric company produces electric bulbs. During production samples of 200 produced bulbs were taken every day for 10 successive days and the defective bulbs found were recorded as under:					
		8, 3, 0, 10, 5, 4, 6, 3, 2, 2					
		Draw control chart for fraction defective. (P-Chart)					
	(2)	For single sampling plan (1000, 50, 2) when $P'=0.01$ , Obtain Pa, ATI, AOQ and ASN. ( $e^{-0.5}=0.6065$ )	7				
(B)	Answer the following: (Any Three)						
	(1)	Define AQL.					
	(2)	$\overline{X}$ and R charts are based on distribution.					
	(3)	Draw an ideal O.C. curve.					
	(4)	Define consumer's risk.					
	(5)	In AOQ curve the maximum value of AOQ is called					

MF-101 4