

Seat No. : _____

DA-112

December-2018

M.Sc., Sem.-I

401 : Physics

**(Quantum Mechanics-I and Mathematical Physics-I)
(New and Old Course)**

Time : 2:30 Hours]

[Max. Marks : 70

1. (A) (i) Discuss perturbation theory for degenerate states. Show that first order correction to energy can be obtained by diagonalizing $m \times n$ matrix. 7
- (ii) Explain stark effect for first excited state of hydrogen atom. Find first order correction to the energy and show that degeneracy is not completely remove when H-atom is placed in a uniform electric field.
- [Hint : Consider matrix elements of perturbed Hamiltonian for which $l = l' = 0, l = l' = 1$ and $m \neq m'$ are zero]. 7

OR

- (i) Show that $W \geq E_0$ and $[\langle H^2 \rangle_\psi - W^2]^{1/2} \geq (W - E_0)$. 7
- (ii) Find out minimum energy of the He-atom using variation method. 7
- (B) Answer any **four** questions : 4
- (1) Show that for the electric dipole when placed in the uniform electric field of intensity E , the term $\frac{1}{2} \alpha E^2$ has unit of energy. Here, α represents polarizability.
- (2) Show that $\nabla_{11} = \nabla_{22}$.
- (3) Write degenerate states for $n = 4$.
- (4) What will be the scalar product corresponding to the eigen state for energy $+ 3eEa$ and $-3eEa$?
- (5) When dipole moment is aligned to uniform electric field, what will be its energy eigen value ?
- (6) What is the numerical value of effective valency of the atom ?

2. (A) (i) Write an equation of propagator $G(\vec{r}, \vec{r}' ; t, 0)$. Find out differential equations for propagator and retarded propagator. 7
- (ii) Show that in Sudden approximation transition probability is directly proportional to time T and matrix element of $H_0 - H$ between final and initial states. 7

OR

- (i) Obtain Bohr-Sommerfeld quantization condition. 7
- (ii) Using Bohr-Sommerfeld quantization condition show that energy of simple harmonic oscillator is given by $E = \hbar\omega \left(n + \frac{1}{2} \right)$. 7
- (B) Answer any **four** questions. 4

- (1) Define Heaviside function.
- (2) What will be the dimension of $S(x)$ in equation $u(x) = A(x) e^{\frac{iS(x)}{\hbar}}$?
- (3) In WKB approximation solution of the Schrödinger equation is expanded in power _____. Complete the statement.
- (4) In non-classical region, kinetic energy of particle is negative. True or false
- (5) If $V(x)$ is slowly varying function, then what will be $\int P(x)dx$?
- (6) What do you mean by classical turning point ?

3. (A) Write the following :
Prove that Laplace transform :

- (i) $L \{ \sin kt \} = \frac{k}{s^2 + k^2}$ and $L \{ t^n \} = \frac{n!}{s^{n+1}}, s > 0, n > -1$ 7
- (ii) Evaluate the inverse Laplace transform, $L^{-1} \left\{ \frac{s^2}{(s^2 + a^2)(s^2 + b^2)} \right\}$. 7

OR

- (i) The motion of a body falling in a resisting medium is given by $m \frac{d^2x(t)}{dt^2} = mg - b \frac{dx(t)}{dt}$, when the retarding force is proportional to the velocity. Find $x(t)$ and $x'(t)$ for the initial conditions $x(0) = x'(0) = 0$. 7
- (ii) Using Laplace transforms, solve the set of equations :
 $y' - 2y + z = 0$, and
 $z' - y - 2z = 0$, with initial conditions: $y(+0) = 1, z(+0) = 0$.
Given : $L \{ e^{-at} \sin bt \} = b / [(s+a)^2 + b^2]$, and
 $L \{ e^{-at} \cos bt \} = (s+a) / [(s+a)^2 + b^2]$

(B) Answer the following : (Any **Three** out of **Five**) **3**

- (1) Write the equation representing linearity operation for Laplace transform.
- (2) Give the definition of Heaviside unit step function.
- (3) Write an expression for the Laplace transform of third derivative of $F(t)$.
- (4) What is the Laplace transform of Dirac delta function
 $L\{\delta(t - t_0)\} = \dots\dots t_0 \geq 0$?
- (5) For $s > 0, L\{1\} = \dots\dots$.

4. (A) Write the answer of the following :

- (i) Prove that every tensor of second rank be resolved into anti-symmetric and symmetric part. **7**
- (ii) Give definition of a group. Explain four properties of a group with relevant examples. Give a group table of order three with different elements A, B, E. **7**

OR

- (i) If A_{ij}, B_{ij}, C_{ij} and D_{ij} tensors of second rank and same type, then prove that $A_{ij} - B_{ij} = D_{ij}$ and $A_{ij} + B_{ij} = C_{ij}$. **7**
- (ii) Define a class. Discuss four properties of a class with relevant examples. **7**
 What do you understand by product of classes ?

(B) Answer any **three** short questions : **3**

1. List the difference between homomorphism and isomorphism.
2. What do you understand by 'conjugate of subgroups' ?
3. Give an example of second rank tensor.
4. Triad has ____ components.
5. 'Tensors can be multiplied by other tensors to form new tensors' This sentence is _____.s (correct, wrong)
