

Seat No. : _____

DC-120

December-2018

M.Sc., Sem.-I

**403 : Mathematics
(Complex Analysis – I)**

Time : 2:30 Hours]

[Max. Marks : 70

1. (a) (i) For any two complex numbers z_1 and z_2 , prove the triangular inequality $|z_1 + z_2| \leq |z_1| + |z_2|$. When does the equality hold ? **14**
- (ii) Prove that the complex numbers z_1, z_2 and the origin form an equilateral triangle only if $z_1^2 + z_2^2 - z_1 z_2 = 0$.

OR

- (b) (i) If $f(z)$ has a finite limit at z_0 , then $f(z)$ is a bounded function in some neighborhood of z_0 . Also verify the inequality $\sqrt{2} |z| \geq |\operatorname{Re} z| + |\operatorname{Im} z|$.
- (ii) If the function $g(z)$ is continuous at $z = z_0$ and the function $f(z)$ is continuous at $g(z_0)$, then the composite function $f[g(z)]$ is continuous at z_0 .

- (c) Attempt any **four** : **4**

- (i) Describe the region $\{z : \operatorname{Im}(z) < |z - 1|^2\}$. State whether the region is a domain.
- (ii) State the definition of connected set.
- (iii) Compute the limit of function $\lim_{z \rightarrow -i} \frac{(iz^3 + 1)}{z^2 + 1}$ if it exists ?
- (iv) If $z_1 = -5$ and $z_2 = -1 + i$, show that $\operatorname{Arg}(z_1 / z_2) = \operatorname{Arg}(z_1) - \operatorname{Arg}(z_2)$.
- (v) Where the function $\operatorname{Ln}(1 + z^2)$ is not continuous on the complex plane ?
- (vi) State the definition of Uniform continuity of a complex function. Is $f(z) = 3z - 2$ is uniformly continuous in the region $|z| \leq 1$?

2. (a) (i) State and prove the sufficient condition for a function to be analytic. 14
- (ii) Show that the function $\text{Ln } z$ is analytic for all z except when $\text{Re } z \leq 0$.

OR

- (b) (i) Let $f(z)$ be an analytic function on a connected open set D . If there are two constants c_1 and $c_2 \in \mathbb{C}$, not all zero, such that $c_1 f(z) + c_2 \overline{f(z)} = 0$ for all $z \in D$, then show that $f(z)$ is a constant on D .
- (ii) If $f(z) = u + iv$ is an analytic function of $z = x + iy$ and $u + v = (x + y)(2 - 4xy + x^2 + y^2)$ then find u, v and the analytic function $f(z)$ in terms of z .

- (c) Attempt any **four** : 4

- (i) Are functions $f(z)$ and $f(\bar{z})$ simultaneously analytic ? If yes, why ?
- (ii) Let $f(z)$ be an analytic function in a domain D . $f(z)$ is constant in D if $f(z)$ vanishes identically in D . Is this statement true ? If yes, why ?
- (iii) Is the function $w = |z|^2$ continuous everywhere but nowhere differentiable except at the origin ? If yes, why ?
- (iv) State Cauchy-Riemann equations in polar form.
- (v) The function $f(z) = z \sec z$ is not analytic at the points $z = \underline{\hspace{2cm}}$
- (vi) If $u = x^2 - y^2$ is harmonic, then $\frac{\partial^2 u}{\partial z \partial \bar{z}} = \underline{\hspace{2cm}}$

3. (a) (i) State and prove the existence of the contour integral. 14
- (ii) Evaluate the integral $I = \int_C (x + y^2 - ixy) dz$, where

$$C : z = z(t) = \begin{cases} t - 2i, & 1 \leq t < 2, \\ 2 - i(4 - t), & 2 < t \leq 3. \end{cases}$$


OR

- (b) (i) Find a simple, piecewise smooth curve in t the form $z = z(t)$, whose trace and direction is given by the square with vertices at $z = 3 \pm 3i$ and $z = -3 \pm 3i$ traversed counter clockwise. Also check whether the curve

$$z(t) = \begin{cases} t + i(1 + t^2), & -2 \leq t \leq 1 \\ 2 - t + i[3 - (t - 2)^2], & 1 < t \leq 4. \end{cases}$$

is closed, simple, smooth or piecewise smooth.

- (ii) Find an upper bound for the absolute value of the integral $I = \int_C (\exp(2z) - z^2) dz$, where C is the contour given in the figure below :



- (c) Attempt any **three** :

3

- (i) The primitive of a function is unique up to an additive constant, why ?
(ii) A simple closed Jordan curve divides the Argand plane into _____ open domains which have the curve as common boundary.
(iii) Compute the length of the curve $z(t) = (1 - i) \exp(-it)$, $0 \leq t \leq \pi/2$.
(iv) Find an upper bound for the absolute value of the integral $I = \int_C \exp((\bar{z})^2) dz$, $C: |z|=2$ where C is traversed in the anticlockwise direction.

4. (a) (i) State and prove Cauchy integral theorem.

14

- (ii) Evaluate the integral $I = \oint_C \frac{\exp(z)}{z^2(z+1)^3} dz$, $C: |z|=2$.

OR

- (b) (i) State and prove Liouville's theorem.
- (ii) Verify that the maximum and minimum modulus theorems hold for the function $f(z) = z^2 + 1$, where C is the circle $|z| = 1$ and D is the domain inside C .

(c) Attempt any **three** :

3

- (i) State Cauchy inequality.
- (ii) If $f(z)$ is an analytic function within and on a simple closed contour C and z_0 is any point inside C , then show that $\oint_C \frac{f(z)}{(z - z_0)^2} dz = \oint_C \frac{f'(z)}{(z - z_0)} dz$.
- (iii) Evaluate the integral $I = \oint_C \frac{\exp(z)}{z^2 + 9} dz$, $C : |z| = 2$.
- (iv) Define Simply connected domain. Can we convert Multiply connected domain into Simply connected domain ? If yes, how ?
-